

Well Child Check and Immunization Schedule

Age	Visit	Immunizations and Tests
Birth	Newborn Exam in Hospital	Newborn Metabolic Screening (NBS) Hepatitis B Nirsevimab Hearing Screen
2-4 days	Well Child Check	Hepatitis B (if not given at birth) Nirsevimab (if not given at birth)
2 weeks	Well Child Check	Hepatitis B (if not given at birth) Nirsevimab (if not given at birth)
2 Months	Well Child Check	Pediarix (DTaP, Hep B, IPV) Pneumococcal Pedvax Hib Nirsevimab (if not given at birth) Rotateq (RV) *Must be given between 6-14 weeks*
4 Months	Well Child Check	Pediarix (DTaP, Hep B, IPV) Pneumococcal Pedvax Hib Nirsevimab (if not given at birth) Rotateq (RV)
6 Months	Well Child Check	Pediarix (DTaP, Hep B, IPV) Pneumococcal Nirsevimab (if not given at birth) Rotateq (RV) *Third dose given before 8 months*
9 Months	Well Child Check	*PPD if indicated
12 Months	Well Child Check	Pneumococcal Pedvax Hib MMR (Measles, Mumps, Rubella) Hemoglobin and Lead Screening
15 Months	Well Child Check	DTaP Hep A Varicella
18 Months	Well Child Check	No shots are required if shots are current Photoscreening
2 Years	Well Child Check	Hep A #2 and Lead Screening
3 Years	Well Child Check	No shots are required if shots are current
4 Years	Well Child Check	Kinrix (DTaP, IPV) MMR (Measles, Mumps and Rubella) Varicella
5 Years	Well Child Check	*MMR, Kinrix, & Varicella (if not given at 4yr visit)
6-12 Years	Well Child Check	Lipid Testing TdaP after 10 years Meningococcal ACYW & Meningococcal B after 11 years HPV (2 shots 6 months apart) after 9 years
13-21 Years	Teen/Young Adult Health Visit	Hemoglobin for menstruating females Tetanus booster when indicated (every 10 years if no injury) Meningococcal ACYW & Meningococcal B booster between 16 & 18 yrs HPV if not given earlier - 2 shots 6 mo apart if started prior to 15 years - 3 shots spaced apart over 6 months if started after 15 years